

**Test 3**

Recitation Section (see back of test): \_\_\_\_\_

- 1) Print your name, test form number (above), and nine-digit student number in the section of the answer card labeled "STUDENT IDENTIFICATION".
- 2) Bubble your test form number (**ABOVE**) in columns 1-3, skip column 4, then bubble in your student number in columns 5-13.
- 3) For each free-response question, show all relevant work supporting your answer. **Clearly box or underline your final answer.** "Correct" answers which are not supported by adequate calculations and/or reasoning will be counted wrong.
- 4) For each multiple-choice question, select the answer most nearly correct, **circle this answer on your test**, and bubble it in on your answer card. **Show all relevant work on your quiz.**
- 5) Be prepared to present your Buzzcard as you turn in your test. Scores will be posted to WebAssign after they have been graded. **Quiz grades become final when the next quiz is given.**
- 6) You may use a simple scientific calculator capable of logarithms, exponentials, and trigonometric functions. **Programmable engineering calculators with text or graphical capabilities are not allowed. Wireless devices are prohibited.**



Your test form is: **533**

The image shows a Scantron answer card with the following details:


- STUDENT IDENTIFICATION:** George P. Burdell
- THRU:** 123
- STUDENT NUMBER:** 900987654
- WRITTEN information for human benefit:** Test Form Number from YOUR test, column 4 is blank, YOUR student number starts in column 5
- Answers:** A box labeled "Your answers" highlights the first four columns (1-4) of the answer grid.
- Grid:** A grid of 25 columns and 10 rows of bubbles for answers.
- SCN-151-10** and copyright information are visible on the right side.

**Our Final Exam will be on Wednesday, July 29!**

The following problem will be hand-graded. Show all your work for this problem. Make no marks and leave no space on your answer card for it.


- [I] (20 points) You are an accident investigator, working the scene of a collision between a sportscar (mass  $m$ ) and a UPS truck (mass  $3m$ ). The truck was initially moving due east into the intersection with a speed  $v_T = 24$  mph (as confirmed by a GPS recorder on the truck). The car was moving due north into the intersection, and the driver claims that he was traveling below the posted speed of 25 mph. Skid marks from the collision indicate that the two vehicles stuck together, sliding in a direction  $\theta = 22.5^\circ$  north of east as they came to a stop.

Determine the actual speed of the sportscar just before the collision. Was the driver really obeying the speed limit, as he claimed?



The following problem will be hand-graded. Show all your work for this problem. Make no marks and leave no space on your answer card for it.

- [III] (20 points) A pan of negligible mass hangs from a spring that is suspended from the ceiling. Initially, the pan hangs a distance  $L$  below the ceiling (*far left*). When a mass  $m$  is gently placed on the pan, the pan hangs at rest a distance  $5L/4$  below the ceiling (*center*). If the same mass  $m$  is dropped from the ceiling onto the pan (*far right*), at what distance  $D$  below the ceiling will the pan momentarily come to a stop (*not shown*)? Express your answer in terms of the parameter  $L$ .




The following problem will be hand-graded. Show all your work for this problem. Make no marks and leave no space on your answer card for it.

- [III] (20 points) A block of mass  $m$  is pushed up a rough ramp at constant speed, by the application of a horizontal pushing force  $P$ . The ramp is inclined at an angle  $\theta = 36.9^\circ$  above the horizontal, and the coefficient of friction for the ramp is  $\mu_k = 0.500$ .

Identify all of the forces acting on the block, and calculate the work done by each force, as the block moves a distance  $D$  up the ramp. **Express each answer in terms of  $M$ ,  $g$ , and/or  $D$ .**

*Hint: start by using the Second Law to find each force as some multiple of  $mg$ !*




*Question value 4 points*

- (1) A hammer is dropped from the roof of a tall building. Ignoring drag, the power delivery by gravity to the hammer will be:
- constant and positive throughout the hammer's descent.
  - initially zero, but increasing uniformly to greater and greater positive values as the hammer descends.
  - initially zero, but decreasing uniformly to greater and greater negative values as the hammer descends.
  - zero, because no power is ever delivered by a conservative force like gravity.
  - constant and negative throughout the hammer's descent.

*Question value 8 points*


- (2) A block hanging from a cord is whirled in a horizontal circle, forming a conical pendulum. What forces act, and what work do they do? (Assume the block is in uniform circular motion.)
- Tension *along* the cord and gravity straight down. Gravity does positive work while tension does an equal amount of negative work.
  - Tension *along* the cord and gravity straight down. Gravity does negative work while tension does an equal amount of positive work.
  - Tension, gravity, and the force of the circular motion. Tension and gravity do zero work, while the force of the motion does positive work.
  - Tension *along* the cord and gravity straight down. Neither force does *any* work.
  - Tension, gravity, and the centripetal force. Since they are all perpendicular to the motion, they all do zero work.



*Question value 8 points*

- (3) In the figure at right, a child of mass  $m$  is riding a rope-swing of length  $L$ . If the child swings back and forth making a maximum angle of  $60^\circ$  from the vertical, what will be the tension in the rope at the very bottom of her (circular!) arc?


- (a)  $mg/2$   
 (b)  $2mg$   
 (c)  $(3/2)mg$   
 (d)  $mg$   
 (e)  $(\sqrt{3}/2)mg$



*Question value 8 points*

- (4) The diagram at right displays a head-on collision between two blocks of unequal mass. What are the two blocks' final velocities, if the collision is perfectly elastic?


- (a)  $\vec{v}_A = \langle -v \rangle$  and  $\vec{v}_B = \langle -v/2 \rangle$   
 (b)  $\vec{v}_A = \langle +v/2 \rangle$  and  $\vec{v}_B = \langle +v \rangle$   
 (c)  $\vec{v}_A = \langle -v \rangle$  and  $\vec{v}_B = \langle +v/2 \rangle$   
 (d)  $\vec{v}_A = \langle -2v \rangle$  and  $\vec{v}_B = \langle +v \rangle$   
 (e)  $\vec{v}_A = \langle -2v \rangle$  and  $\vec{v}_B = \langle -v \rangle$



*Question value 8 points*

- (5) The graph at right displays the horizontal force on a tennis ball by a wall, during the time interval that the ball is rebounding off the wall. What is the magnitude of the average force exerted on the ball by the wall?

- (a) 4.8 N
- (b) 4.0 N
- (c) 1.6 N
- (d) 8.0 N
- (e) 2.4 N



*Question value 4 points*

- (6) A car is traveling due west at some speed  $v_0$ . It then turns right, and ends up traveling due north with the same speed  $v_0$ . What impulse (if any) does the ground deliver to the car during this process?

- (a) There is no impulse at all during this process, because the car's speed didn't change.
- (b) There is a northeastward-directed impulse on the car by the ground.
- (c) There is a southeastward-directed impulse on the car by the ground.
- (d) There is a northwestward-directed impulse on the car by the ground.
- (e) There is a southwestward-directed impulse on the car by the ground.

## PHYS 2211M Recitation TA and Room Assignments

Tests will be returned in recitation, in the week *after* the test. In order to ensure that you receive your test back as soon as possible, please enter your recitation section from the table above on the front of this test.

	<b>Room 123</b>
<b>TUESDAY</b>	
2:20–3:10 pm	M04 Kothari, Kartik
3:20–4:10 pm	M03 Kothari, Kartik
4:20–5:10 pm	M01 Ge, Luwei
<b>WEDNESDAY</b>	
2:20–3:10 pm	
3:20–4:10 pm	M05 Shi, Chao
4:20–5:10 pm	M02 Shi, Chao
<b>THURSDAY</b>	
2:20–3:10 pm	M08 Ge, Luwei
3:20–4:10 pm	

---